

**BIOLOGY AND MORPHOLOGY OF THE BANANA MOTH,
OPOGONA SACCHARI (BOJER), AND ITS INTRODUCTION
INTO FLORIDA (LEPIDOPTERA: TINEIDAE)**

DONALD R. DAVIS AND JORGE E. PEÑA

(DRD) Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560; (JEP) Institute of Food and Agricultural Sciences, University of Florida, Homestead, Florida 33031.

Abstract.—The general distribution and recent introduction into Florida of the banana moth, *Opogona sacchari* (Bojer) is reviewed. Currently, the principal damage caused by this species in Florida consists of larval stem boring in certain nursery stock and ornamental palms in particular. The biology of the species is summarized and all stages of the insect are described, supplemented by numerous illustrations.

Key Words: Lepidoptera, Tineidae, *Opogona sacchari*, banana moth, moth biology, immature stages

The banana moth, *Opogona sacchari* (Bojer), is known from several tropical-subtropical humid regions around the world but has yet to be reported from the Indo-Australian region. Originally reported from the Mascarene Islands in the Indian Ocean (Bojer 1856, Walker 1863, Butler 1876), the species was later discovered in Africa (Vari and Kroon 1986) as well as islands near the African continent (Walker 1875, Durrant 1925) and Europe (Ciampolini 1973, D'Aguilar and Martinez 1982, Mourikis and Vassilaina-Alexopoulon 1983). More recently the species has spread to South America (Cintra 1975, Giannotti et al. 1977) and the West Indies (Alam 1984). The earliest evidence we have of this species in the New World is represented by adult specimens in the Smithsonian Institution (USNM) that were collected in 1970 from Aragua, Venezuela on potato.

Specimens received for identification from B. Kumashiro of the Hawaiian Department of Agriculture just prior to publication indicate that *O. sacchari* is now established in Hawaii. The species has been reared from

rotting coconut tree tops (central whorl of leaves) from Kaneohe, Oahu and from an unidentified palm at Kohala, Hawaii.

Within the last few years *O. sacchari* has become established on various nursery stock in southern Florida, particularly in Dade and Palm Beach Counties (Heppner, Peña, and Glenn 1987). Nursery stock particularly affected in Florida include corn plant or casse (*Dracena fragrans* (L.) Ker-Gaus, variety *massangeana*) and bamboo palms (*Chamaedorea* sp.) as well as Hawaiian good luck plant (*Cordyline terminalis* (L.) Kunth) and aralias (*Polyscias* sp.). Although sugar cane is a major host, *O. sacchari* has not yet been reported on that plant in the United States.

The appearance of this new pest in the United States has necessitated a careful examination of all developmental stages and a report of its biology pertinent to its current significance as a pest of nursery stock in Florida.

Opogona sacchari (Bojer)

Alucita sacchari Bojer 1856: 21, pl. 5, figs. 1-10.

Opogona sacchari (Bojer).—Vinson, 1938: 56 (synonym of *Opogona subcervinella*).—Viette, 1957: 145; 1958: 4.—Ciampolini, 1973: 221.—Cintra, 1975: 223.—Giannotti et al., 1977: 209.—Declercq and Van Luchene, 1977: 499.—Zimmerman, 1978: 386.—Cintra et al., 1978: 3.—Pigatii, 1978: 21.—Pigatii et al., 1979: 61.—Veenenbos, 1981: 235.—D'Aguilar and Martinez, 1982: 28.—Rotundo and Tremley, 1982: 123.—Suplicy and Sampaio, 1982: 174.—Bennett and Alam, 1985: 41.—Heppner et al., 1987: 1.

Tinea subcervinella Walker, 1863: 477.

Opogona subcervinella (Walker).—Walsingham, 1907: 713; 1919: 259.—Meyrick, 1930: 321.—Vinson, 1938: 56.—Viette, 1951: 339; 1957: 145 (synonym of *Opogona sacchari*); 1958: 4.—Paulian and Viette, 1955: 147.—Box, 1953: 34.—Davis, 1978: 14; 1984: 22.—Vari and Kroon, 1986: 84, 156.

Hieroxestis subcervinella (Walker).—Meyrick, 1910: 375; 1911: 298.—Cockerell, 1923: 247.—Meyrick, 1924: 556.—Durrant, 1925: 12.—Oldham, 1928: 147.—Rebel, 1939: 63; 1949: 56.—Jannone, 1966: 24.

Gelechia sanctaehelenae Walker, 1875: 192.—Durrant, 1925: 12 (synonym of *Hieroxestis subcervinella*).

Euplocamus sanctaehelenae (Walker).—Wollaston, 1879: 417.

Hieroxestis sanctaehelenae (Walker).—Durrant, 1923: xvii.

Gelechia ligniferella Walker, 1875: 192.—Durrant, 1925: 12 (synonym of *Hieroxestis subcervinella*).


Laverna plumipes Butler, 1876: 409.—Walsingham, 1907: 713 (synonym of *Opogona subcervinella*); 1919: 259.

Adult (Figs. 1, 2).—Length of forewing: ♂, 7.3–11.2 mm; ♀, 9–12.5 mm. Moderately large, generally dark grayish brown moths with a small, black subapical spot on forewing near apex of discal cell and a similar one at basal third on anal fold; male slightly

paler in color with faint longitudinal streaks of light brown or buff.

Head: Vestiture smooth except for a pair of bilateral tufts of erect, pale brown to cream, piliform scales between antennal bases; scales of frons broad, cream colored; vertex covered by a dense row of broad cream scales that curve over frons; caudal portion of vertex with broad, grayish brown scales; occiput light brown with a scattered arch or median patch of dark fuscous scales. Antenna 103–125 segmented, approximately 0.8 the length of forewing; scape cream dorsally, with scattered fuscous scales ventrally; flagellum uniformly covered with small but moderately broad, cream to buff scales arranged in a single row, completely encircling each flagellomere (Fig. 12); sensilla chaetica (with spiral grooves) and trichodea (longitudinal grooves) relatively dense and randomly scattered over each flagellomere; a few pair of sensilla coeloconica located near distal margin (Figs. 13, 14). Pilifers moderately developed with setae nearly meeting at midline (Fig. 9). Labrum densely covered with microtrichia. Mandible vestigial but exceeding pilifer setae in length. Maxillary palpus 5-segmented, elongate, exceeding length of relatively short haustellum; dorsal and lateral surfaces densely covered with cream scales; venter naked with dense sensilla; apex of fifth segment with a slender basiconic sensillum and another slightly smaller one at subapex (Figs. 15, 16). Haustellum with a series of shallow plates over basal half; largest plates with a pair of short sensilla basiconica (Fig. 10). Labial palpus upcurved, smoothly covered with cream scales except for a lateral, subapical series of 3–4 cream bristles and a ventral subapical tuft of 6–8 bristles; third segment with an elongate, narrow sensory pit located just beyond middle (Figs. 17, 18).

Thorax: Pronotum light brown with heavy to sparse scattering of fuscous scales and usually with a small to large median patch of fuscous on anterior margin and on tegula. Venter uniformly shiny cream. Forewing




Figs. 1-4. *Opogona sacchari*. 1, Adult male, length of forewing 9.5 mm. 2, Adult female, length of forewing 12.5 mm. 3, Larvae, maximum length 30 mm. 4, Cocoon with pupal exuvium, cocoon length 12.5 mm.

light brown in female with an almost equal amount of scattered dark brown scales; male with dark brown scales more concentrated into longitudinal streaks, particularly evident along upper and lower margins of discal cell; a small, dark fuscous subapical spot near apex of cell and a more elongate fuscous spot along anal fold at basal third in both sexes; male retinaculum a relatively elongate flap from ventral subcostal margin, with apical margin rolled under (Figs. 24, 25). Hindwing light yellowish brown with slender scales in male, more shiny gray and with slightly broader scales in female; male with elongate hair pencil from dorsal base (Figs. 28-30); ultrastructure of pencil sex scales as illustrated (Figs. 31-33); male frenulum a single stout bristle (Fig. 26); female

with usually five stout frenular bristles (Figs. 27). Foreleg mostly cream; coxa with a small amount of grayish fuscous suffusion at base; dorsal surfaces of tibia and tarsus heavily suffused with grayish fuscous; normal pectinated epiphysis present (Figs. 19, 20). Pretarsus of all legs normal, with relatively broad unguitactor plate bearing 7-10 ranks of scutes in a single transverse row (Figs. 21-23). Midleg uniformly cream except for slight grayish fuscous suffusion over dorsal surface of tarsus. Hindleg similar to midleg in color except with less grayish fuscous on tarsus and with dense, elongate piliform setae from tibia.

Abdomen: Terga mostly grayish brown with caudal margins and pleura buff to cream colored; venter of female uniformly cream,



5



6




7




8


Figs. 5-8. Damage to *Dracaena fragrans* by *Opogona sacchari* larvae (see arrows). 5, Healthy stem. 6-8, Typical feeding damage to *Dracaena fragrans* canes.




Figs. 9–14. *Opogona sacchari*, adult morphology. 9, Labrum, anterior view (100 μm). 10, Base of haustellum (38 μm). 11, Haustellum, food canal (15 μm). 12, Antenna, scale pattern (50 μm). 13, Antenna, sensilla (60 μm). 14, Sensillum coeloconicum of antenna (4.3 μm). (Scale lengths in parenthesis; bar scale for all photographs = Fig. 9.)




Figs. 15–20. *Opogona sacchari*, adult morphology. 15, Maxillary palpus, apical (fifth) segment (100 μm). 16, Apex of Fig. 15 (23.1 μm). 17, Labial palpus, apical (third) segment (176 μm). 18, Detail of sensory pit in Fig. 17 (see arrow) (25 μm). 19, Epiphysis (120 μm). 20, Detail of pecten on epiphysis (27 μm). (Scale lengths in parentheses; bar scale for all photographs = Fig. 15.)



Figs. 21–26. *Opogona sacchari*, adult morphology. 21, Pretarsus of midleg, lateral view (20 μm). 22, Ventral view (50 μm). 23, Unguitractor plate of pretarsus (7.5 μm). 24, Subcostal retinaculum of male forewing (0.27 mm). 25, Distal-lateral view of Fig. 24 (176 μm). 26, Male frenulum (38 μm). (Scale lengths in parentheses; bar scale for all photographs = Fig. 21.)



Figs. 27–32. *Opogona sacchari*, wing morphology. 27, Female frenulum (250 μm). 28, Hair pencil of male hindwing, dorsal wing (1 mm). 29, Base of male hair pencil (176 μm). 30, Raised scale sockets of hair pencil (30 μm). 31, Surface ultrastructure of a single hair pencil scale (3.8 μm). 32, Detail of Fig. 31 (1.2 μm). (Scale lengths in parentheses; bar scale for all photographs = Fig. 27.)



Figs. 33–38. *Opogona sacchari*, adult, egg, and larval morphology. 33, Cross section of hair pencil scale (Fig. 31) ($3.8\ \mu\text{m}$). 34, Egg, lateral view ($150\ \mu\text{m}$). 35, Micropyle ($136\ \mu\text{m}$). 36, Detail of Fig. 35 ($75\ \mu\text{m}$). 37, Detail of central disk (Fig. 36) ($17.6\ \mu\text{m}$). 38, Head of larva, anterior view ($60\ \mu\text{m}$). (Scale lengths in parentheses; bar scale for all photographs = Fig. 33.)

grayish brown in male; both sexes with a lower lateral series of five dark fuscous spots on A3-7.

Male genitalia (Figs. 79-82): Uncus deeply divided into two large widely separated lobes arising beneath tegumen and bearing numerous stout elongate setae on their mesal surfaces. Tegumen a relatively broad ring dorsally. Vinculum tapering to a broad, relatively truncate saccus. Valva prominently divided into a large, elongate, rounded apical lobe and a much smaller, acute cucullar lobe. Aedoeagus a relatively small, straight, slender tube without cornuti; phallobase much larger and greatly inflated.

Female genitalia (Figs. 83, 84): Tertiary apophyses present in A10. Genital plate moderately divided into a pair of rounded lobes. Corpus bursae with a single large sagittate signum bearing elongate anterior arms.

Egg (Figs. 34-37).—Length, 0.5-0.55 mm; diameter, 0.38 mm; oval in shape, round in cross section; color light yellow at oviposition, gradually becoming yellowish brown prior to eclosion. Surface irregularly pitted. Micropyle consisting of a single, centrally positioned opening with small radiating grooves forming an enlarged, reticulate pattern of low ridges over entire end of egg; reticulations mostly 5-6 sided.

Larva (Figs. 3, 38-66).—Length of largest larva 30 mm, maximum diameter 3 mm. Body generally white with dark brown plates and pinacula.


Head: Maximum width 2.5 mm. Color reddish brown, becoming darker anteriorly and over frons. Chaetotaxy as illustrated. Stemmata rudimentary, consisting of a pair of widely separated, clear lenses, probably corresponding to stemmata 1 and 5; the most anterior (5) located ventral to antennal socket; the most posterior (1) midway between A3 and S2. Mandible with five cusps. Spinneret long and slender with a minute orifice. Labial palpus 2-segmented, elongate and slender, equalling or slightly exceeding length of spinneret; apical seta one-third the

length of entire palpus. Apex of mentum with a pair of minute secondary labial setae.

Thorax: Pronotum and spiracular plate dark reddish brown, lighter in color along margins; spiracle together with L setae on same plate. Pinacula over entire body relatively large, dark brown and very distinct on whitish integument. Meso- and meta-thorax with L2 arising on a separate pinacula from L1 and 3. MSD1 and 2 of similar length and reduced. Legs well developed; tarsal claw elongate; basal lobe actually bilobed and with a minute conical seta from inner angle (Figs. 51, 52).

Abdomen: Whitish in color with brownish pinacula. L2 on separate pinacula from spiracle. Prolegs well developed on A3-6 and 10; crochets A3-6 uniordinal, uniserial, and arranged in a complete ellipse composed of approximately 43-45 hooks; a scattered band of much smaller, numerous spines encircling apex of planta (Fig. 53); crochets on A10 with 20-22 hooks and a dense, scattered band of much smaller spines along anterior edge of planta (Figs. 55, 56).


Pupa (Figs. 67, 68).—Length, 9-12.8 mm; maximum width, 2-3.5 mm. Color usually light brown ventrally and dark reddish brown dorsally; wing cases becoming darker with maturity; frontal process, cremaster, and adjacent areas extremely dark, often black. Head with frontal process (cocoon cutter) moderately developed, with a broad, triangular apex (Figs. 68, 78); labrum with a pair of lateral setae. Anterolateral margins of mesonotum with a pair of perforated bands composed of minute, raised slit openings (Figs. 69-71). Forewings extending to middle of A5. Hindlegs to A6. A simple, anterior row of short, stout, dorsal spines present on segments A4-8; tabulation of spines as follow: A4 = 56-70, A5 = 53-72, A6 = 57-71, A7 = 44-50, A8 = 8-18. Last pair (A8) of spiracles on raised, swollen bases (Figs. 74-76). Cremaster consists of a large pair of stout hooks arising dorsally from




Figs. 39–44. *Opogona sacchari*, larval morphology. 39, Head, dorsal view (0.6 mm). 40, Labrum, dorsal view (176 μ m). 41, Head, lateral view (0.43 μ m). 42, Stemmal area (136 μ m). 43, Head, ventral view (0.5 mm). 44, Maxilla and labium (150 μ m). (Scale lengths in parentheses; bar scale for all photographs = Fig. 39.)




Figs. 45–50. *Opogona sacchari*, larval morphology. 45, Maxilla (60 μm). 46, Apex of maxillary palpus (6 μm). 47, Apex of maxilla (5 μm). 48, Apex of antenna (30 μm). 49, Apex of antenna (27 μm). 50, Ventral view of pro- and mesothorax (0.6 mm). (Scale lengths in parentheses; bar scale for all photographs = Fig. 45.)




Figs. 51–56. *Opogona sacchari*, larval morphology. 51, Pretarsal claw of prothorax (50 μm). 52, Detail of axial lobes and seta of Fig. 51 (6 μm). 53, Proleg of A3, \uparrow anterior, \rightarrow meson (136 μm). 54, Caudal view last abdominal segment, A10 (0.5 mm). 55, Ventral view, A9, 10 (0.5 mm). 56, Anal proleg, A10 (150 μm). (Scale lengths in parentheses; bar scale for all photographs = Fig. 51.)




Figs. 57-64. *Opogona sacchari*, larval chaetotaxy. 57, Body segments T1-2, A1, 6, 8-9. 58, Head, dorsal view (0.5 mm). 59, Ventral view. 60, Segments A8-10, dorsal view. 61, Head, lateral view. 62, Mandible. 63, Labrum, dorsal view (0.25 mm). 64, Ventral view. (Scale lengths in parentheses.)



Figs. 65-70. *Opogona sacchari*, larval and pupal morphology. 65, Larval segments A8-10, dorsal view (0.6 mm). 66, Lateral view of Fig. 65 (0.6 mm). 67, Pupa, lateral view of head (0.6 mm). 68, Ventral view of Fig. 67 (0.6 mm). 69, Dorsal view of Fig. 67, perforated band (see arrow) (0.6 mm). 70, Detail of dorsal perforated band (136 μ m). (Scale lengths in parentheses; bar scale for all photographs = Fig. 65.)



Figs. 71-76. *Opogona sacchari*, pupal morphology. 71, Detail of openings in perforated band (see Fig. 69) (15 μ m). 72, Dorsum of A4-5 (0.75 mm). 73, Dorsal spines of A4 (100 μ m). 74, Dorsal view of A7-10 (0.5 mm). 75, Caudal view of A8-10 (0.5 mm). 76, Lateral view of A8-10 (0.5 mm). (Scale lengths in parentheses; bar scale for all photographs = Fig. 71.)



Figs. 77, 78. *Opogona sacchari*, pupa. 77, Ventral view (2 mm). 78, Dorsal view. (Scale lengths in parentheses.)

A10 (Figs. 74–76); a much smaller pair of conical tubercles also present ventrally.

Pupation occurs in a cocoon (Fig. 4) of white silk usually covered with dark frass and plant debris, 14–18 mm long, 3–4 mm in diameter. Normally, the cocoon is constructed somewhere in or near the feeding site.

Types.—Syntypes ♂, ♀, deposition unknown (*sacchari*); Holotype, BMNH (*subcervinella*); syntypes, ♂, ♀, BMNH (*sanctaelenae*); holotype, sex unstated, BMNH (*ligniferella*); syntype(s)?, sex and number of specimens unstated, BMNH (*plumipes*).

Type localities.—Mascarene Islands: Mauritius (*sacchari*, *subcervinella*), St. Helena (*sanctaelenae*, *ligniferella*); Mascarene Islands: Rodriguez (*plumipes*).


Hosts.—46 plant hosts reported (see Table 1).

Distribution (maps 1, 2).—A widely ranging, partly pantropical species, early reported from several circum-African islands (Canary Islands, St. Helena, Mauritius,

Rodriguez and the Seychelles), as well as South Africa; frequently intercepted in several European countries (Belgium, France, Great Britain, Greece, and the Netherlands). More recently, *O. sacchari* has been introduced into South America and the West Indies, including Bermuda, and is now established in nurseries over much of southern Florida.

Discussion.—By its relatively large size and predominantly brown color, *Opogona sacchari* is easily distinguished from nearly all other members of this large complex. Its affinities are with *O. omoscopia* (Meyrick) as suggested by their similar male genitalia and shared presence of a dorsal hair pencil in the hindwings of the male. Present evidence suggests that this complex of obviously related species warrants recognition under and resurrection of the currently synonymized genus *Hieroxestis*. Hopefully this and similar generic uncertainties can be resolved following a revision of the large cosmopolitan genus *Opogona*. The larva of *O. sacchari* may be recognized by the presence of two stemmata, separation of the spiracle from the pinaculum bearing L2 on the first eight abdominal segments, by the large number of crochets (A3–6 = 43–45, A10 = 20–22), and by complete encirclement of the abdominal planta by a band of small, secondary spines. In *O. omoscopia*, a pantropical species recently introduced into California (Davis 1978), only one anterior stemma has been observed, the first eight abdominal spiracles are united with L2 on a common pinaculum, the crochets are fewer in number, and the abdominal planta A3–6 have spines only along the anterior margin. The pupae of these two species are also similar but may be distinguished by the raised spiracle on A8 and larger cremaster spines of *O. sacchari*.

A peculiar structure of unknown function in the pupa of this species deserves further comment. It consists of a bilateral pair of perforated bands curving around the anterolateral margins of the mesonotum (Figs.



Figs. 79-84. Adult genitalia. 79, Male, ventral view (0.25 mm). 80, Lateral view. 81, Lateral view of valva. 82, Aedeagus, lateral view. 83, Female, ventral view (1 mm). 84, Detail of signum in Fig. 83. (Scale lengths in parentheses.)

69-71). The bands are narrow and elongate and under high magnification can be observed to comprise raised, slitlike openings. The senior author has not noted this in other species, perhaps because the bands were simply overlooked. Possibly their function is merely to weaken the cuticle, thereby facilitating rupture of the pupal shell during

eclosion. The band was observed to have been expanded or further separated in most pupal exuviae examined, although the major breaks in this region of the pupal shell during ecdysis did not occur through the perforated bands but were located anterior to the pronotum and down the mid-dorsal line of the pro- and mesonotum.

Table 1. Plant hosts of *Opogona sacchari*.

Plant Species	Reference	Country
Agavaceae		
<i>Cordylone terminalis</i> (L.) Kunth	Heppner et al. 1987	United States
<i>Dracaena fragrans</i> (L.)	Declercq and Van Luchene 1977	Belgium
<i>Dracaena fragrans</i> (L.) Ker-Gaus, "var. <i>massangeana</i> "	Heppner et al. 1987	United States
<i>Dracaena marginata</i> Lam.	Heppner et al. 1987	United States
<i>Dracaena reflexa</i> Lam.	Heppner et al. 1987	United States
<i>Yucca elephantipes</i> Regel	Heppner et al. 1987	United States
<i>Yucca</i> sp.	Heppner et al. 1987	United States
Araceae		
<i>Colocasia esculenta</i> Schott.	Cintra 1975	Brazil
<i>Philodendron scandens</i> Lindl.	Süss 1974	Italy
Araliaceae		
<i>Polyscias fruticosa</i> (L.) Harms	Heppner et al. 1987	United States
<i>Polyscias fruticosa</i> (L.) Harms, "elegans"	Heppner et al. 1987	United States
Asteraceae		
<i>Dahlia</i> sp.	Cintra 1975	Brazil
Bromeliaceae		
<i>Aechmea fasciata</i> (Lindl.) Baker	Süss 1974	Italy
<i>Aechmea fasciata</i> (Lindl.) Baker "Variegata"	Süss 1974	Italy
<i>Guzmania lingulata</i> var. \times <i>magnifica</i> [Hort.]	Süss 1974	Italy
<i>Nidularium tricolor</i> [species name unknown, possibly = <i>Neoregelea</i> 'Perfecta Tricolor,' a cultivar]	Süss 1974	Italy
Caricaceae		
<i>Carica papaya</i> L.	Viette 1951	Madagascar
Convolvulaceae		
<i>Ipomoea batatas</i> Lam.	USNM (new record)	Peru
Cycadaceae		
<i>Cycas revoluta</i> Thunberg	Heppner et al. 1987	United States
Dioscoreaceae		
<i>Dioscorea</i> sp.	Heppner et al. 1987	United States
Gesneriaceae		
<i>Gloxinia</i> sp.	Süss 1974	Italy
<i>Saintpaulia</i> sp.	Süss 1974	Italy
Iridaceae		
<i>Gladiolus</i> sp.	Cintra 1975	Brazil
Leguminosae		
<i>Albizia julibrissin</i> Durazz.	Heppner et al. 1987	United States
<i>Enterolobium</i> sp.	Heppner et al. 1987	United States
<i>Erythrina variegata</i> L.	FSCA (new record)	United States
Liliaceae		
<i>Sansevieria laurantii</i> Wildem.	Süss 1974	Italy
<i>Sansevieria trifasciata laurentii</i> Wildem.	Declercq and Van Luchene 1977	Belgium
Malvaceae		
<i>Hibiscus</i> sp.	Moreton 1974	Great Britain

Table 1. Continued.

Plant Species	Reference	Country
Marantaceae		
<i>Maranta leuconeura massangeana</i> Schum.	Süss 1974	Italy
<i>Stromanthe sanguinea</i> Sonder	Süss 1974	Italy
Moraceae		
<i>Ficus elastica</i> (H. A. Siebrecht)	Moreton 1974	Great Britain
Musaceae		
<i>Musa cavendishii</i> Paxt	Oldham 1928	Canary Islands
<i>Musa paradisiaca</i> L.	Heppner et al. 1987	United States
<i>Musa sapientum</i> L.	Oldham 1928	Canary Islands
<i>Strelitzia</i> sp.	Zandvoort 1972	Netherlands
Orchidaceae		
"Orchids"	Heppner et al. 1987	United States
Palmae		
<i>Arecastrum</i> sp.	Heppner et al. 1987	United States
<i>Bactris</i> [= <i>Guilielma</i>] <i>gasipaes</i> HBK	Heppner et al. 1987	United States
<i>Chamaedorea elegans</i> Mart.	Heppner et al. 1987	United States
<i>Chamaedorea erumpens</i> H. E. Moore	Heppner et al. 1987	United States
<i>Chamaedorea seifrizii</i> Burret	Heppner et al. 1987	United States
Poaceae		
"Bamboo"	Oldham 1928	Canary Islands
<i>Saccharum officinarum</i> L.	Bojer 1856	Mauritius
<i>Zea mays</i> L.	Oldham 1928	Canary Islands
Solanaceae		
<i>Capsicum</i> sp.	Süss 1974	Italy
<i>Solanum melongena</i> L. var. <i>cuculentum</i> Nees	Süss 1974	Italy
<i>Solanum tuberosum</i> L.	Oldham 1928	Canary Islands
Verbenaceae		
<i>Clerodendrum</i> sp.	Heppner et al. 1987	United States

BIOLOGY


The larvae of most species of *Opogona* for which we have information are detritus feeders and rarely feed on living plant tissue. *Opogona sacchari* thus departs from the norm and can be a serious pest of banana, maize, potato, sweet potato, sugar cane, and certain greenhouse crops (Alam 1984, Durrant 1925, Oldham 1928, Süss 1975, Veenbos 1981). A total of seven instars are indicated, based on head capsule measurements of larvae reared on artificial diet (Table 2). All measurements were based on pooled data for both sexes. The discreteness of the last two instars as interpreted from head capsule measurements presumably

would be better had males only or females only been used. The SD values do not overlap as it is, so likely the conclusion reached as to instar number is correct. More precise data, however, might have shown indication of variation in instar numbers, which seems likely in a generalist feeder of this sort.

Oldham (1928) observed that *O. sacchari* larvae feed on nearly all parts of the banana plant except the roots and leaf blades. Larvae accepted leaves as a food in rearings but normally avoided this part of the plant in nature. The most serious damage occurred in the banana inflorescence. Larvae seldom feed exposed but burrow into the substra-



Map 1. Reported occurrence of *Opogona sacchari*.



Map 2. Distribution of *Opogona sacchari* in Florida (after Heppner et al. 1987).

tum. Their presence is usually indicated by the accumulation of frass and other debris entangled in larval silk over the surface of the injury. The larvae are voracious feeders and construct long meandering galleries through the injury site.

Reports of larval damage on sugar cane has varied markedly. On Barbados, Alam (1984) noted extensive damage to live sugar cane, exceeding that caused by *Diatraea saccharalis* (F.). The young larvae feed under the leaf sheaths and, as they mature, penetrate the stalks and destroy the cane tissue. Infested stalks are hollowed out and gradually filled with larval frass. As in the case on banana, pupation occurs at the feeding site inside the stalks. At the infestation sites studied by J. E. Jones (in litt.), also on Barbados, *O. sacchari* was most abundant in dead or dying stumps and dead canes, suggesting they moved in following an infestation by *Diatraea*. Jones also found that the high incidence of *Opogona* on green canes was associated with a high infestation

Table 2. Larval head capsule width for 7 instar groupings of *Opogona sacchari* (Bojer).


Instar	N	Range (mm)	Mean \pm SD (mm)	Ratio of Increase
1	26	0.16–0.24	0.18 \pm 0.03	
2	20	0.26–0.32	0.31 \pm 0.02	1.72
3	27	0.36–0.56	0.49 \pm 0.01	1.51
4	20	0.58–0.88	0.91 \pm 0.02	1.85
5	20	0.91–1.16	1.02 \pm 0.04	1.12
6	20	1.20–1.44	1.42 \pm 0.04	1.39
7	32	1.52–2.56	2.17 \pm 0.04	1.42
Average				1.52

of *Diatraea*, again suggesting that *O. sacchari* invades burrows previously created by *Diatraea*. Thus, the observations by Jones agree most closely with those originally reported by Bojer (1856).

OBSERVATIONS IN FLORIDA

A laboratory colony of *O. sacchari* was initiated in 1985 with larvae and pupae collected from infested *Dracaena* canes found at a nursery in Homestead, Florida. Pupae were held individually in 36 ml diet cups until eclosion. Emerging adults were sexed and 17 pairs were placed in 11 cm \times 14.5 cm plastic oviposition chambers along with folded filter paper. Adults were provided honey and water. Pairs were held until the female died, males were replaced as needed. The preoviposition period, fecundity and longevity of females and the time required for egg to hatch were determined. First instars were gathered from the oviposition chambers and placed on one of 3 artificial diets in an attempt to rear this species in the laboratory. The 3 diets used were: velvet bean caterpillar diet (VBC), sugar cane borer diet, and the elm spanworm diet (Fedde 1974). All rearing and diet tests were conducted in a rearing room with a photoperiod of 12:12 (L:D) (0600 to 1800 photophase), temp 24 \pm 2°C, and RH 65–70%. Larvae were collected at different days from the diet, and preserved in a 70% alcohol for later measurement of head capsule width. Larvae

Table 3. Mating activity of moths in rearing chamber.



were checked daily for onset of pupation and pupal development time was recorded for 175 pupae.

Emerging adults were also placed in 53 × 53 × 50 cm screen cages with *Dracaena* and *Chamaedorea* potted plants. Number of eggs and oviposition site on each plant species was inspected daily.

RESULTS AND DISCUSSION

Based on laboratory observations, development of an *Opogona* generation required 50–70 days. More eggs were present on unexpanded leaves and stems than on expanded leaves. Eggs are laid singly or in groups up to 328. They are light yellow at oviposition, turn a dark yellow color ca. 2 days later, and finally yellowish brown prior to eclosion. Eggs hatched in 7.02 ± 0.02 days at 24°C in the laboratory. The preoviposition period for newly emerged females in the laboratory was 1.66 ± 0.14 days at

25°C. Female longevity ($\bar{x} \pm SD$, $n = 20$) in the laboratory was 9–17 days with a mean of 11.45 ± 0.72 days. Sixty first instar larvae were placed individually on each of the 3 diets tested. Survivorship on the artificial diet was low with the exception of velvet bean caterpillar diet. The highest percentage survivorship, 83%, was on VBC diet.

Experiments were conducted to monitor *O. sacchari* sexual activity throughout the night. Sets ($n = 30$) of 2–3-day-old virgin males and females were placed in petri dishes (9 cm in diameter) with water and honey as source of food. Each set was placed in an environmental chamber (LD 12:12, $24 \pm 2^\circ\text{C}$, 75–80% RH). The experiment was replicated 4 times. Diel activity was monitored hourly from 6:00 pm to 7:00 am. The period of activity occurred between 1:00 am to 4:00 am. No sexual activity was observed before 1:00 am or after 4:00 am. Through this experiment it became apparent that the op-

timal response period for *O. sacchari* occurs 3 h before the end of scotophase (darkness) (Table 3).

The pupal stadium of larvae placed on VBC diet lasted 12.53 ± 0.33 days. The average weight of pupae was 0.043 g. Table 2 lists a 7 instar model which best fits the data according to Dyar's rule (Dyar 1890). As noted previously, these measurements are based on pooled data for unsexed larvae.

Typical damage of *O. sacchari* on *Dracaena* is characterized by removal of the bark and phloem. Cuttings of *Dracaena* having *Opogona* larvae show exterior debris and frass (Figs. 7, 8) deposits, and have internal feeding damage on dead and living portions of the cortex, pith, roots and leaves (Fig. 6). Damage is not evident 4–6 weeks after infestation. Typical damage to *Chamaedorea* palms can be observed 2–3 weeks after infestation. Each larva feeds at the base and roots of *Chamaedorea*, and frass accumulates at the plant base from feeding into roots and petioles. In palms the leaf blades of the growing point become bleached and necrotic.

ACKNOWLEDGMENTS

The senior author (Davis) is responsible for the systematic and morphological portions of this paper and the biological observations in Florida are by the junior author (Peña).

We are indebted to Vichai Malikul and Young Sohn of the Department of Entomology, Smithsonian Institution, for the line drawings and to Susann Braden and Brian Kahn of the Smithsonian SEM Lab and Victor Kranz of the Smithsonian Photographic Laboratory for photographic assistance. The final draft of the manuscript was prepared by Callie Sullivan. We wish to thank Joël Minet and Pierre Viette of the Museum of National d'Histoire Naturelle (Paris), John Heppner of the Florida State Collection of Arthropods (Gainesville, Florida), and Kevin Tuck of the British Museum (Natural

History, London) for providing information essential to our report. We also appreciate the comments by two anonymous reviewers of the manuscript.

LITERATURE CITED

- Alam, M. M. 1984. New insect pests of sugarcane in Barbados. Barbados Sugar Technologists' Association, Second Annual Conference, pp. 1–6.
- Bennett, F. D. and M. M. Alam. 1985. An annotated check-list of the insect and allied terrestrial arthropods of Barbados. Caribbean Agricultural Research and Development Institute, pp. 1–81.
- Bojer, W. 1856. Report of the committee on the "cane borer." Pp. 1–46, 5 pls. H. Plaidean Government Printer, Port Louis, Mauritius.
- Box, H. E. 1953. List of sugarcane insects. Pp. 1–101. Commonwealth Institute of Entomology, London.
- Butler, A. G. 1876. Preliminary notice of new species of Lepidoptera from Rodriguez. *Annals and Magazine of Natural History*, 44th Series 17(101): 407–409.
- Ciampolini, M. 1973. *Opogona sacchari* Bojer dannosa a piante ornamentali. *Notiziario sulle Molluschi delle Piante*, Nos. 88–89, pp. 221–225. Società SIPCAM, Milan, Italy.
- Cintra, A. F. 1975. *Opogona* sp. nova praga da bananicultura em São Paulo. *Biológico* 41(8): 223–231.
- Cintra, A. F., P. R. Almeida, I. Myazaki, and H. S. Neves. 1978. Fumigação de banana com brometo de metila para controle de lagartas de *Opogona sacchari* (Bojer, 1856). *Biológico* 44(1): 3–10.
- Cockerell, T. D. A. 1923. The Lepidoptera of the Madeira Islands. *The Entomologist* 56: 247.
- D'Aguilar, J. and M. Martinez. 1982. *Opogona sacchari* (Bojer) présent dans les cultures sous serres en France (Lep. Tineidae). *Bulletin de la Société Entomologique de France* 87(1/2): 28–30.
- Davis, D. R. 1978. The North American moths of the genera *Phaeoses*, *Opogona*, and *Oinophia*, with a discussion of their supergeneric affinities (Lepidoptera: Tineidae). *Smithsonian Contributions to Zoology* 282: 1–39, 128 figs.
- . 1984. Tineidae, pp. 4–5, 19–24. In Heppner, J. B., ed., *Atlas of Neotropical Lepidoptera*, vol 2 (checklist), part 1 (Micropterigoidea to Immoidea). Dr. W. Junk, the Hague.
- Declercq, R. and I. Van Luchene. 1977. Voorkomen en bestrijding van de rupsen van *Opogona sacchari* Boj. in *Dracena fragrans* en *Sansevieria trifasciata* 'Laurentii.' *Verbondsnieuws voor de Belgische Sierteelt* 21(15): 499, 501, 1 pl.

- Durrant, J. H. 1923. A moth from Saint Helena in Britain. *Proceedings of the Entomological Society of London*, p. xvii.
- . 1925. The banana moth, *Hieroxestis subcervinella* Wlk. *Entomologist's Monthly Magazine* 61: 12–13.
- Dyar, H. G. 1890. The number of molts of Lepidopterous larva. *Psyche* 5: 420–422.
- Fedde, Y. H. 1974. Rearing the elm spanworm on artificial diet: A preliminary report. USDA Forest Service, Ashville, N. C., SE-204, pp. 1–5.
- Giannoti, O., B. S. Oliveira, T. Toneda, and D. Fell. 1977. Observacoes gerais sobre o desenvolvimento e comportamento sexual do Lepidoptera *Opogona sacchari* (Bojer, 1856) em laboratório. *Arquivos do Instituto Biológico* 44(4): 209–212.
- Heppner, J. B., J. E. Peña, and H. Glenn. 1987. The banana moth, *Opogona sacchari* (Bojer) (Lepidoptera: Tineidae), in Florida. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. *Entomology Circular No. 293*, pp. 1–4.
- Jannone, G. 1966. Risultati di alcuni controlli fitosanitari su vegetalie prodotti vegetali esteri nel Porto di Genova. *Rivista di Agricoltura subtropicale e Tropicale* 60(1–3): 5–34.
- Meyrick, E. 1910. Descriptions of Microlepidoptera from Mauritius and the Chagos Isles. *Transactions of the Entomological Society of London*, pp. 366–377.
- . 1911. Tortricina and Tineina of the Percy Sladen Trust Expedition to the Indian Ocean in 1905. *Transactions of the Linnaean Society of London, Zoology, 2nd series* 14: 263–307.
- . 1924. Microlepidoptera of Rodriguez. *Transactions of the Entomological Society of London*, pp. 544–557.
- . 1930. Microlepidoptera of Mauritius. *Transactions of the Entomological Society of London* 78: 309–323.
- Moreton, B. D. 1974. New or uncommon plant diseases and pests. *Plant Pathology* 23: 163–164.
- Mourikis, P. A. and P. Vassilaina-Alexopoulou. 1983. Data on the biology of the *Opogona sacchari* (Bojer 1856), a new pest for ornamental plants in Greece. *Anales de L'institute Phytopathologique Benaki* 13(1): 59–64.
- Oldham, J. N. 1928. *Hieroxestis subcervinella* Wlk., an enemy of the banana in the Canary Islands. *Bulletin of Entomological Research* 19(pt. 2): 147–166, pls. 4–5.
- Paulin, R. and P. Viette. 1955. Essai d'un catalogue biologique des Lepidopteres heteroceres de Tananarive. *Memoirs de L'Institut Scientifique de Madagascar, ser. E* 6: 141–281.
- Pigatti, A. 1978. Seleção de inseticidas em laboratório para o controle da 'traca' da banana—*Opogona sacchari* (Bojer, 1856) (= *O. subcervinella* Walker, 1863)—Lepidoptera: Lyonetiidae. *Biológico* 44(1): 21–23.
- Pigatti, A., P. Almeyda, A. Cinta, and D. Oliveira. 1979. Ensaio com inseticidas aplicados por via liquida para o controle da "traca da banana"—*Opogona sacchari* (Bojer, 1856) (= *O. subcervinella*, Walker, 1863) Lepidoptera: Lyonetiidae. *Biológico* 45(3/4): 61–68.
- Rebel, H. 1939. Achten Beitrag zur Lepidopterenfauna der Kanaren. *Annalen des Naturhistorischen Museums in Wien* 49: 43–68.
- . 1949. Anhang. Eine Lepidopteren-Ausbente von Madeira. *Die Lepidopteren fauna des Azorischen Archipels* 8(1): 50–57.
- Rotundo, G. and E. Tremlay. 1982. Preliminary observations on the sex pheromone of *Opogona sacchari* (Bojer). *Bollettino del laboratorio di Entomologia Agraria Filippo Silvestri* 39: 123–132.
- Suplicy Fo., N. and A. S. Sampaio. 1982. Pragas da Bananeira. *Biológico* 48(7): 169–182.
- Süss, L. 1975. *Opogona sacchari* (Bojer) (Lepidoptera, Lyonetiidae), a new pest of ornamental plants in greenhouses. *Bollettino di Zoologia di Entomologia Agraria dell Universita* 12: 1–28.
- Vari, L. and D. Kroon. 1986. Southern African Lepidoptera, a series of cross-referenced indices. 198 pp. The Lepidopterists Society of South African and the Transvaal Museum, Pretoria.
- Veenenbos, J. A. V. 1981. *Opogona sacchari*, a pest risk from imports of ornamental plants of tropical origin. *Bulletin, Organisation Européene et Méditerranéenne pour la Protection des Plantes* 11(3): 235–237.
- Viette, P. 1951. Quelques Microlepidopteres de Madagascar. *Memoires de L'Institut Scientifique de Madagascar, série A* 5(2): 335–347.
- . 1957. Lepidopteres (excepte les tordeuses) et les Geometrides. *Memoires de L'Institut Scientifique de Madagascar, série E* 8: 137–226.
- . 1958. Lepidoptères Tinéides (s.l.) et Pyrales. *Societas Scientiarum Fennica. Commentationes Biologicae*, XVII, 8 17(8): 1–12.
- Vinson, J. 1938. Catalogue of the Lepidoptera of the Mascarene Islands. *Mauritius Institute Bulletin* 1 (pt. 4): 1–69.
- Walker, F. 1863. List of the specimens of Lepidopterous insects in the collections of the British Museum 28: 287–561.
- . 1875. Fam. Gelechiidae [sic], p. 192. *In* Melliss, J. C., ed., *St. Helena: A Physical, Historical, and Topographical Description of the Island, Including its Geology, Fauna, Flora, and Meteorology*. xiv + 426 + pp., pls. 1–56. L. Reece & Co., London.

- Walsingham, Lord (Thomas de Gray). 1907. Microlepidoptera. In Sharp, D., ed., Fauna Hawaiiensis, or the Zoology of the Sandwich (Hawaiian) Isles 1(pt. 5): 469-759, pls. 10-22. Cambridge University Press.
- . 1919. Madeiran Tineina (Lepidoptera). The Entomologist's Monthly Magazine, second series 21(46): 257-259.
- Wollaston, E. 1879. Notes on the Lepidoptera of St. Helena, with descriptions of new species. Annals and Magazine of Natural History, fifth series 3: 219-233, 329-343, 415-441.
- Zandvoort, R. 1972. Bestrijding van *Opogona subcervinella* bij *Strelitzia*. Gewasbescherming 3(4): 95-96.
- Zimmerman, E. C. 1978. Insects of Hawaii. Microlepidoptera 9: 1-1876, figs. 1-1355. Honolulu: University of Hawaii Press.